arXiv:1207.3238v1 [math.ST] 13 Jul 2012

Using Moving Average Method To Estimate Entropy In Testing Exponentiality For Type-II Censored Data

A. Kohansal and S. Rezakhah*

July 16, 2012

Abstract

In this paper, we introduce a modified test statistic, by applying moving average method and present a new cdf estimator to estimate the joint entropy of the type-II censored data. We also establish a goodness of fit test statistic based on the Kullback-Leibler information for the type-II censored data, and compare its performance with the leading test statistic. A Monte Carlo simulation study shows that the proposed test statistic has greater power than the leading test statistic against the alternatives with monotone increasing hazard functions.

Keywords: Entropy, Monte Carlo simulation, Kullback-Leibler distance, moving average method, Hazard function.

Mathematics Subject Classification: 62G10, 62G30.

1 Introduction

Suppose that a random variable X has a distribution function $F(x)$, with a continuous density function $f(x)$. The differential entropy $H(X)$ of the random variable X is defined by [22] to be

$$H(X) = -\int_{-\infty}^{\infty} f(x) \ln f(x) dx.$$

The nonparametric estimation of $H(X)$ has been discussed by many authors including Vasicek [25], Theil [24], Dudewicz and van der Meulen [6], Ebrahimi et al. [8] and Bowman [4]. While Vasicek [25] and Ebrahimi et al. [8] directly obtained the

---

*Faculty of Mathematics and Computer Science, Amirkabir University of Technology, Tehran, Iran. Email: rezakhah@aut.ac.ir, ak_kohansal@aut.ac.ir

nonparametric entropy estimators, Theil [24], Dudewicz and van der Meulen [6] and Bowman [4] derived the nonparametric entropy estimators from the nonparametric distribution functions.
The entropy difference $H(X) - H(Y)$ has been considered in establishing goodness of fit tests for the class of the maximum entropy distributions [7], [12].

The Kullback-Leibler (KL) information in favor of X against Y with the density functions $g(x)$ and $f(x)$ is defined to be

$$I(X;Y) = \int_{-\infty}^{\infty} g(x) \ln \frac{g(x)}{f(x)} dx.$$

Because $I(X;Y)$ has the property that $I(X;Y) \geq 0$, and the equality holds if and only if $f = g$, the estimate of the KL information has been also considered as a goodness of fit test statistic by some authors including [1], [9]. It has been shown in the aforementioned papers that the test statistics based on the KL information perform very well for exponentiality [9], and s-normality [25] in terms of powers compared with some leading test statistics for complete samples.

From the censored data point of view, some authors studied the problem of goodness of fit test and discussed some test statistics based on a censored data sample. In the case of type-II censored data, if $X_{(1:n)}, \cdots, X_{(r:n)}$ be the first r order statistics of a random sample of size n from a distribution with cdf $F(x)$, then the joint entropy of these data is defined as:

$$H_{1 \cdots r:n} = -\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{x_{2:n}} f_{1 \cdots r:n} \ln f_{1 \cdots r:n} dx_{(1:n)} dx_{(r:n)}, \tag{1.1}$$

where $f_{1 \cdots r:n}$ is the joint pdf of $X_{(1:n)}, \cdots, X_{(r:n)}$.
However $H_{1 \cdots r:n}$, is an r multiple integral, and we need to simplify the multiple integral. The simple calculation of the entropy of single and consecutive order statistics has been studied in [19], [26].

[18] showed that the multiple integral $H_{1 \cdots r:n}$ can be simplified to a single integral as:

$$H_{1 \cdots r:n} = -(\ln n + \cdots + \ln(n-r+1)) + n\bar{H}_{1 \cdots r:n},$$

where

$$\bar{H}_{1 \cdots r:n} = \frac{r}{n} - \int_{-\infty}^{\infty} (1 - F_{r:n-1}(x)) f(x) \ln h(x) dx,$$

where h is the corresponding hazard function.
He also showed that:

$$\bar{H}_{1 \cdots r:n} = -E\left(\int_0^{U_{r:n-1}} \ln\left(\frac{d}{dp} F^{-1}(p)\right) dp\right) - E((1 - U_{r:n-1}) \ln(1 - U_{r:n-1})),$$

and proposed the following estimator of $\bar{H}_{1\cdots r:n}$ as:

$$\bar{H}_{m,n,r} = \frac{1}{n}\sum_{i=1}^{r} \ln\left(\frac{n}{2m}(x_{(i+m:n)} - x_{(i-m:n)})\right) - \left(1-\frac{r}{n}\right)\ln\left(1-\frac{r}{n}\right),$$

where m is a positive integer less than or equal to $\frac{r}{2}$, which is called window size and $x_{(i-m:n)} = x_{(1:n)}$ for $i \leq m$ and $x_{(i+m:n)} = x_{(r:n)}$ for $i \geq r-m$. Also [27] suggested another estimator of $\bar{H}_{1\cdots r:n}$ based on the new cdf estimator.

In the case of progressively censored data, first Balakrishnan et al. [3] studied the testing exponentiality based on KL information with progressively type-II censored data. Goodness of fit test based on KL information for progressively type-II censored data can be found in Habibi Rad et al. [13].

The rest of the article is arranged as follows: In Section 2, we provide a new test statistic: first, we explain the simple moving average method and then, we present our test statistic. The statistical properties of the proposed test statistic are obtained in section 3. Implementation of the testing exponentiality based on type-II censored data is evaluated in Section 4 and in this section it is shown that, for some types of alternatives, the proposed test achieves higher power than the leading test statistic.

2 Presentation of new test statistic

2.1 Simple moving average method

A simple moving average (SMA) method is the unweighted mean of the previous n datum points. The SMA method are common tools in technical analysis [16] and [10].

Definition 2.1 *Given a sequence a_i, $i = 1\cdots N$, an n-moving average is a new sequence S_i, $i = 1\cdots N-n+1$ defined from the a_i by taking the arithmetic mean of subsequences of n terms:*

$$S_i = \frac{1}{n}\sum_{j=i}^{i+n-1} a_j,$$

So the sequences S_n giving n-moving averages are

$$S_2 = \frac{1}{2}(a_1 + a_2, a_2 + a_3, \cdots, a_{n-1} + a_n),$$

$$S_3 = \frac{1}{3}(a_1 + a_2 + a_3, a_2 + a_3 + a_4, \cdots, a_{n-2} + a_{n-1} + a_n),$$

and so on.

The order selected n depends on the type of movement of interest, such as short, intermediate, or long term. An SMA can also be disproportionately influenced by old datum points dropping out or new data coming in.

One characteristic of the SMA is that if the data have an uneven path, then applying the SMA will eliminate abrupt variation and cause the smooth path. In the next subsection, we use the definition and this characteristics of SMA method and present a new test statistic.

2.2 New test statistic

Some non-parametric estimators of (1.1) have been proposed by [18] and [27]. [27] expressed (1.1) in the form $H_{1\cdots r:n} = -\ln \frac{n!}{(n-r)!} + n\bar{H}_{1\cdots r:n}$, where

$$\bar{H}_{1\cdots r:n} = -E\left(\int_0^{U_{(r:n-1)}} \ln(\frac{dF^{-1}(p)}{dp})dp\right) - E\left(\left(1-U_{(r:n-1)}\right)\ln(1-U_{(r:n-1)})\right),$$

with approximating $\bar{H}_{1\cdots r:n}$ by

$$-\int_0^{\frac{r}{n}} \ln(\frac{dF^{-1}(p)}{dp})dp - (1-\frac{r}{n})\ln(1-\frac{r}{n}), \tag{2.2}$$

in which $\frac{r}{n} = E(U_{(r:n-1)})$, they provided its estimator as

$$\bar{H}^*_{m,n,r} = \frac{1}{n}\sum_{i=1}^{r} \ln\left(\frac{x_{(i+m:n)} - x_{(i-m:n)}}{\hat{F}_n(x_{(i+m:n)}) - \hat{F}_n(x_{(i-m:n)})}\right) - (1-\frac{r}{n})\ln(1-\frac{r}{n}) \tag{2.3}$$

where

$$\hat{F}_n(x_{(i:n)}) = \frac{r-1}{r(n+1)}\left(i + \frac{1}{r-1} + \frac{x_{(i:n)} - x_{(i-1:n)}}{x_{(i+1:n)} - x_{(i-1:n)}}\right), \quad i = 1, \cdots, r,$$

in which m is a positive integer less than or equal to $\frac{r}{2}$, which is called window size and $x_{(i-m:n)} = x_{(1:n)}$ for $i \leq m$ and $x_{(i+m:n)} = x_{(r:n)}$ for $i \geq r-m$.

$F^{-1}(p)$ as a function of quantiles in (2.2) is the sample path of order statistics, but it usually is not smooth, so relation (2.3) can not provided a good estimator for (2.2). Then we propose to imply SMA method of proper order, say k, to smooth this sample path and re-consider (2.3) for the new variables $y_1, \cdots, y_r$. So by considering such smoothing method, we introduce our new test statistic.

For a null distribution function $f^0(x;\theta)$, the KL distance for the type-II censored data is:

$$I_{1\cdots r:n}(f, f^0) = \int_{-\infty}^{\infty} f_{1\cdots r:n}(x;\theta)\ln\frac{f_{1\cdots r:n}(x;\theta)}{f^0_{1\cdots r:n}(x;\theta)}dx.$$

Then the KL distance can be approximated with

$$I_{1\cdots r:n}(f, f^0) = -n\bar{H}_{1\cdots r:n} - \sum_{i=1}^{r} \ln f^0(x_i;\theta) - (n-r)\ln(1 - F^0(x_r;\theta)).$$

Thus the test statistic based on $I_{1\cdots r:n}(f;f^0)/n$ and $y_1, \cdots, y_r$ can be written as:

$$R_{m,n,r} = -\bar{H}^{+}_{m,n,r} + \frac{r}{n}\left(\ln\left(\frac{1}{r}\left(\sum_{i=1}^{r} y_{(i:n)} + (n-r)y_{(r:n)}\right)\right) + 1\right), \tag{2.4}$$

where $\hat{\theta} = \frac{1}{r}\left(\sum_{i=1}^{r} y_{(i:n)} + (n-r)y_{(r:n)}\right)$ is maximum likelihood estimator of θ and

$$\bar{H}^{+}_{m,n,r} = \frac{1}{n}\sum_{i=1}^{r} \ln \frac{y_{(i+m:n)} - y_{(i-m:n)}}{\hat{F}_n(y_{(i+m:n)}) - \hat{F}_n(y_{(i-m:n)})} - \left(1 - \frac{r}{n}\right)\ln\left(1 - \frac{r}{n}\right), \tag{2.5}$$

and

$$\hat{F}_n(y_{(i:n)}) = \frac{r-1}{r(n+1)}\left(i + \frac{1}{r-1} + \frac{y_{(i:n)} - y_{(i-1:n)}}{y_{(i+1:n)} - y_{(i-1:n)}}\right), \quad i = 1, \cdots, r,$$

in which

$$\begin{aligned}
&y_1 = \tfrac{x_{(1:r)} + \cdots + x_{(k:r)}}{k},\\
&y_2 = \tfrac{x_{(2:r)} + \cdots + x_{(k+1:r)}}{k},\\
&\quad\vdots\\
&y_{r-k+1} = \tfrac{x_{(r-k+1:r)} + \cdots + x_{(r:r)}}{k-1},\\
&y_{r-k+2} = \tfrac{x_{(r-k+2:r)} + \cdots + x_{(r:r)}}{k-2},\\
&\quad\vdots\\
&y_{r-1} = \tfrac{x_{(r-1:r)} + x_{(r:r)}}{2},\\
&y_r = x_{(r:r)}.
\end{aligned} \tag{2.6}$$

For $y < y_{(1:n)}$, $\hat{F}_n(y)$ is less than $\frac{1}{n+1}$ and for $y > y_{(r:n)}$, $\hat{F}_n(y)$ is more than $\frac{r}{n+1}$. In the followings we assume that m in (2.4) and (2.5) is a positive integer less than or equal to $\frac{r}{2} + k$, which is called window size and $y_{(i-m:n)} = y_{(1:n)}$ for $i \leq m$ and $y_{(i+m:n)} = y_{(r:n)}$ for $i \geq r - m$.
Since I is non-negative and is zero if and only if $f = f^0$, a.e., we reject the null hypothesis for large values of $R_{m,n,r}$.

Example 2.1 *For the explanation of the our proposed method, we simulate 30 samples form the exponential distribution, consider their order statistics and censor 5 of them from right then we plot these 25 points in Figure 1.*



Figure 1: Sample path of order statistics of 30 samples from the exponential distribution.

Smoothing of the sample path of order statistics, and using estimator (2.3) for new variables could provide a better estimation for (2.2). So we smooth sample path of order statistics by SMA, and provide a better estimation with greater power. We choose SMA of order 3 and define new variables as:

$$\begin{aligned}
&y_1 = \tfrac{x_{(1:25)}+x_{(2:25)}+x_{(3:25)}}{3},\\
&y_2 = \tfrac{x_{(2:25)}+x_{(3:25)}+x_{(4:25)}}{3},\\
&\ \vdots\\
&y_{23} = \tfrac{x_{(23:25)}+x_{(24:25)}+x_{(25:25)}}{3},\\
&y_{24} = \tfrac{x_{(24:25)}+x_{(25:25)}}{2},\\
&y_{25} = x_{(25:25)},
\end{aligned}$$

and plot the smoothed path of new variables in Figure 2.

Figure 2 shows that the new sample path is smoother than the sample path of the original order statistics. If we choose SMA with order 5, it defines new variables as:



Figure 2: Smoothed sample path of order statistics from order 3 at SMA.

$$y_1 = \frac{x_{(1:25)}+x_{(2:25)}+x_{(3:25)}+x_{(4:25)}+x_{(5:25)}}{5},$$
$$y_2 = \frac{x_{(2:25)}+x_{(3:25)}+x_{(4:25)}+x_{(5:25)}+x_{(6:25)}}{5},$$
$$\vdots$$
$$y_{21} = \frac{x_{(21:25)}+x_{(22:25)}+x_{(23:25)}+x_{(24:25)}+x_{(25:25)}}{5},$$
$$y_{22} = \frac{x_{(22:25)}+x_{(23:25)}+x_{(24:25)}+x_{(25:25)}}{4},$$
$$y_{23} = \frac{x_{(23:25)}+x_{(24:25)}+x_{(25:25)}}{3},$$
$$y_{24} = \frac{x_{(24:25)}+x_{(25:25)}}{2},$$
$$y_{25} = x_{(25:25)}.$$

So we find an smoother sample path by plotting theses new variables in Figure 3.

This Figure shows that even though smoothing sample path of order statistics by SMA of order 3 is not as smooth as using SMA with order 5, but resulting powers, which is demonstrated in section 4, are the same up to two digit of decimals. So without loss of generality, we just consider SMA of order $k = 3$ *in (2.6).*

3 Statistical properties of proposed test statistic

In this section, we consider the some statistical properties of the proposed test statistic in (2.4). The following theorem which states that the scale of the random variable X has no effect on the accuracy of $\bar{H}^{+}_{m,n,r}$ in estimating $H_{1\cdots r:n}$, can easily



Figure 3: Smoothed sample path of order statistics from order 5 at SMA.

be proved by following the line of argument in [9].

Theorem 3.1 *Let* $H^{Y}_{1\cdots r:n}$ *and* $H^{W}_{1\cdots r:n}$ *denote entropies of the distribution of continuous random variables* Y *and* W*, respectively, and* $W = kY$*, where* $k > 0$*. Then:*

- $E(\bar{H}^{+\,W}_{m,n,r}) = E(\bar{H}^{+\,Y}_{m,n,r}) + \frac{r}{n}\ln k$,
- $Var(\bar{H}^{+\,W}_{m,n,r}) = E(\bar{H}^{+\,Y}_{m,n,r})$,
- $MSE(\bar{H}^{+\,W}_{m,n,r}) = E(\bar{H}^{+\,Y}_{m,n,r})$,

where the superscript Y *and* W *refer to the corresponding distribution.*

Proof: It is easy to see that:

$$\hat{F}^{Y}_{n}(w_{(i:n)}) = \frac{r-1}{r(n+1)}\left(i + \frac{1}{r-1} + \frac{w_{(i:n)} - w_{(i-1:n)}}{w_{(i+1:n)} - w_{(i-1:n)}}\right) = \hat{F}^{W}_{n}(y_{(i:n)})$$

for $i = 1, \cdots, r$. So we have:

$$\bar{H}^{+\,W}_{m,n,r} = \frac{1}{n}\sum_{i=1}^{r} \ln \frac{ky_{(i+m:n)} - ky_{(i-m:n)}}{\hat{F}_n(ky_{(i+m:n)}) - \hat{F}_n(ky_{(i-m:n)})} -$$

$$\left(1 - \frac{r}{n}\right)\ln\left(1 - \frac{r}{n}\right) = \frac{r}{n}\ln k + \bar{H}^{+\,Y}_{m,n,r}.$$

■

Theorem 3.2 *$|\bar{H}^{+}_{m,n,r} - \bar{H}_{1\cdots r:n}| \to 0$ in probability as $n, m \to 0$ and $\frac{m}{n} \to 0$.*

Proof: Since [23]

$$\hat{f}(x_i) \to f(x_i) \text{ in probability,}$$

and [20] $f(y_i) \to f(x_i)$ in MSE $\Rightarrow f(y_i) \to f(x_i)$ in probability, consequently

$$\hat{f}(y_i) \to \hat{f}(x_i) \text{ in probability.} \tag{3.7}$$

On the other hand, we have $x_{(i+m:n)}, x_{(i-m:n)}$ belong to an interval in which $f(x)$ is positive and continuous, then there exists a value $x'_i \in (x_{(i+m:n)}, x_{(i-m:n)})$ such that

$$\frac{\hat{F}_n(x_{(i+m:n)}) - \hat{F}_n(x_{(i-m:n)})}{x_{(i+m:n)} - x_{(i-m:n)}} = \hat{f}(x'_i)$$

consequently from (3.7)

$$\frac{\hat{F}_n(y_{(i+m:n)}) - \hat{F}_n(y_{(i-m:n)})}{y_{(i+m:n)} - y_{(i-m:n)}} = \hat{f}(y'_i) \to$$

$$\hat{f}(x'_i) = \frac{\hat{F}_n(x_{(i+m:n)}) - \hat{F}_n(x_{(i-m:n)})}{x_{(i+m:n)} - x_{(i-m:n)}} \text{ in probability.}$$

and $\bar{H}^{+}_{m,n,r} \to \bar{H}^{*}_{m,n,r}$ in probability.
Since [27] $\bar{H}^{*}_{m,n,r} \to H_{1\cdots r:n}$ in probability, the proof is complete. ■

4 Implementation of the testing exponentiality based on type-II censored data

Suppose that we are interested in a goodness of fit test for

$$\begin{cases} H_0 : f^0(x) = \frac{1}{\theta}\exp(-\frac{x}{\theta}), \\ H_1 : f^0(x) \neq \frac{1}{\theta}\exp(-\frac{x}{\theta}), \end{cases}$$

where θ is unknown.

There are lots of test statistics for exponentiality concerning uncensored data [2], [11], [14], [15] and [17], but only some of them can be extended to the censored data [5], [21], [18] and [27]. About the exponentiality test, against the one class of alternatives [18] showed that the power of his test statistic is greater than [5] and [21] also [27] showed that the power of their test statistic is greater than [18], So, we compare the power of our test statistic with that of the test proposed by [27] based on the test statistic:

$$A_{m,n,r} = -\bar{H}^{*}_{m,n,r} + \frac{r}{n}\left(\ln\left(\frac{1}{r}\left(\sum_{i=1}^{r} x_{(i:n)} + (n-r)x_{(r:n)}\right)\right) + 1\right),$$

Table 1: Values of the window size m

r	m
4-5	5
6-7	6
8-9	7
10-11	8
12-13	9
r-(r+1)	m^* (for even r)

where $\bar{H}^*_{m,n,r}$ is explained in (2.2).

Because the sampling distributions of $A_{m,n,r}$ and $R_{m,n,r}$ are intractable, we determine the percentage points using 10000 Monte Carlo samples from an exponential distribution. In determining the window size m which depend on $n,\ ,r$ and the α, we define the optimal window size m to be one which gives minimum critical points in the sense of [9]. However, we find from the simulated percentage points, the optimal window size m. In view of these results, our recommended values of m for different r are listed in Table 1, where $m^* = r/2 + 3$. The critical values of $R_{m,n,r}$ corresponding to the optimum values of m, are given in Table 2.

Because the proposed test statistic essentially is related to the hazard function, we consider the alternatives according to the type of hazard functions as follows:

- Monotone decreasing hazard: Chi-square with degree of freedom 1 (A1), Gamma with shape parameter 0.5 (A2), Weibull with shape parameter 0.5 and 0.8 (A3 and A4 respectively).
- Monotone increasing hazard: Uniform (B1), Weibull with shape parameter 2 (B2), Gamma with shape parameter 1.5, 2 (B3, B4 respectively), Chi-square with degree of freedom 3, 4 (B5, B6 respectively), Beta with shape parameters 1 and 2, 2 and 1 (B7, B8 respectively).
- Non-monotone hazard: Log normal with shape parameter 0.6, 1.0, 1.2 (C1, C2, C3 respectively), Beta with shape parameters 0.5 and 1.0 (C4).

We consider here the sample size to be 30, and draw conclusions. We made 10000 Monte Carlo simulations for $n = 30$ to estimate the powers of our proposed test statistic and that of the competing test statistic, for $\alpha = 0.1$. The simulation results are summarized in Figures 4-6 and Tables 3-5. In these tables $R_{m,n,r}$ and $A_{m,n,r}$ denote the test statistics of the proposed test and the test proposed by [27] respectively.

Table 2: The critical values of $R_{m,n,r}$

n	r	$\alpha = 0.1$	$\alpha = 0.05$	$\alpha = 0.025$
10	5	0.3445	0.4253	0.5087
	6	0.3251	0.4128	0.5026
	7	0.3136	0.4099	0.4929
	8	0.3104	0.4046	0.4915
	9	0.3101	0.4038	0.4902
20	10	0.1474	0.1913	0.2310
	11	0.1426	0.1830	0.2229
	12	0.1408	0.1828	0.2197
	13	0.1369	0.1805	0.2181
	14	0.1322	0.1773	0.2168
	15	0.1294	0.1740	0.2160
	16	0.1281	0.1742	0.2161
	17	0.1280	0.1739	0.2073
	18	0.1268	0.1649	0.2072
	19	0.1200	0.1620	0.1988
30	15	0.0865	0.1168	0.1500
	16	0.0859	0.1152	0.1430
	17	0.0843	0.1124	0.1383
	18	0.0829	0.1090	0.1380
	19	0.0824	0.1083	0.1355
	20	0.0815	0.1079	0.1311
	21	0.0806	0.1043	0.1294
	22	0.0777	0.1038	0.1281
	23	0.0759	0.1027	0.1280
	24	0.0741	0.0988	0.1275
	25	0.0699	0.0976	0.1265
	26	0.0662	0.0977	0.1219
	27	0.0635	0.0910	0.1200

Table 3: Power comparison (distributions having decreasing hazard function)

Distributions	A1		A2		A3		A4	
r	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$
15	0.0202	0.0018	0.0209	0.0019	0.0562	0.0005	0.0291	0.0221
16	0.0092	0.0021	0.0089	0.0018	0.0313	0.0002	0.0246	0.0210
17	0.0122	0.0016	0.0126	0.0007	0.0462	0.0001	0.0187	0.0194
18	0.0064	0.0009	0.0067	0.0013	0.0269	0.0001	0.0190	0.0184
19	0.0078	0.0006	0.0071	0.0010	0.0408	0.0001	0.0185	0.0167
20	0.0023	0.0007	0.0029	0.0008	0.0189	0.0001	0.0169	0.0152
21	0.0037	0.0008	0.0036	0.0010	0.0250	0.0001	0.0157	0.0151
22	0.0014	0.0008	0.0013	0.0006	0.0118	0.0001	0.0147	0.0127
23	0.0012	0.0009	0.0013	0.0006	0.0189	0.0000	0.0138	0.0126
24	0.0006	0.0009	0.0008	0.0006	0.0047	0.0000	0.0092	0.0106
25	0.0007	0.0005	0.0008	0.0004	0.0093	0.0001	0.0101	0.0111
26	0.0005	0.0005	0.0007	0.0005	0.0019	0.0000	0.1011	0.0093
27	0.0008	0.0005	0.0008	0.0007	0.0020	0.0001	0.1186	0.0086



Figure 4: Power comparison: monotone decreasing hazard alternative at 10% when the sample size is 30.

Table 4: Power comparison (distributions having increasing hazard function)

Distributions	B1		B2		B3		B4	
r	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$
15	0.2190	0.2622	0.8890	0.9217	0.3974	0.4347	0.7065	0.7440
16	0.2383	0.2848	0.9118	0.9420	0.4106	0.4516	0.7254	0.7653
17	0.2633	0.3222	0.9306	0.9534	0.4318	0.4840	0.7510	0.8011
18	0.3148	0.3504	0.9533	0.9689	0.4537	0.4958	0.7774	0.8278
19	0.3369	0.3905	0.9645	0.9762	0.4721	0.4968	0.8034	0.8452
20	0.3867	0.4491	0.9766	0.9831	0.4898	0.5046	0.8272	0.8529
21	0.4418	0.4915	0.9790	0.9889	0.4955	0.5201	0.8380	0.8581
22	0.5002	0.5600	0.9880	0.9912	0.5127	0.5402	0.8573	0.8638
23	0.5799	0.6234	0.9927	0.9950	0.5394	0.5518	0.8690	0.8727
24	0.6313	0.6987	0.9936	0.9963	0.5327	0.5558	0.8688	0.8728
25	0.7297	0.7698	0.9971	0.9974	0.5583	0.5672	0.8823	0.8870
26	0.7894	0.8433	0.9980	0.9983	0.5503	0.5759	0.8961	0.8981
27	0.8691	0.8978	0.9981	0.9990	0.5491	0.5831	0.8873	0.9039

Distributions	B5		B6		B7		B8	
r	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$
15	0.3980	0.4362	0.7072	0.7455	0.1483	0.1754	0.9556	0.9750
16	0.4113	0.4508	0.7242	0.7689	0.1505	0.1822	0.9680	0.9854
17	0.4290	0.4712	0.7529	0.7951	0.1671	0.1956	0.9816	0.9903
18	0.4504	0.4887	0.7896	0.8133	0.1891	0.2120	0.9897	0.9955
19	0.4741	0.4909	0.7959	0.8300	0.1907	0.2311	0.9930	0.9982
20	0.4943	0.5031	0.8184	0.8549	0.2152	0.2494	0.9976	0.9986
21	0.5045	0.5131	0.8375	0.8581	0.2408	0.2744	0.9985	0.9994
22	0.5195	0.5293	0.8537	0.8602	0.2598	0.2962	0.9993	0.9999
23	0.5340	0.5423	0.8726	0.8788	0.2955	0.3233	0.9999	1.0000
24	0.5328	0.5509	0.8746	0.8809	0.3313	0.3656	1.0000	1.0000
25	0.5626	0.5719	0.8795	0.8891	0.3752	0.4132	1.0000	1.0000
26	0.5610	0.5691	0.8971	0.8999	0.4463	0.5693	1.0000	1.0000
27	0.5583	0.5701	0.8991	0.9001	0.4957	0.5221	1.0000	1.0000



Figure 5: Power comparison: monotone increasing hazard alternative at 10% when the sample size is 30.

Table 5: Power comparison (distributions having non-monotone hazard function)

Distributions	C1		C2		C3		C4	
r	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$	$A_{m,n,r}$	$R_{m,n,r}$
15	0.9990	0.9993	0.6505	0.6083	0.3451	0.3039	0.0142	0.0033
16	0.9992	0.9994	0.6518	0.6058	0.3334	0.2833	0.0056	0.0030
17	0.9994	0.9994	0.6477	0.6006	0.3255	0.2763	0.0085	0.0029
18	0.9991	0.9993	0.6518	0.5867	0.3068	0.2481	0.0035	0.0033
19	0.9994	0.9996	0.6452	0.5722	0.2971	0.2291	0.0055	0.0032
20	0.9998	0.9996	0.6267	0.5417	0.2659	0.2047	0.0029	0.0039
21	0.9997	0.9999	0.6117	0.5105	0.2547	0.1829	0.0033	0.0038
22	0.9999	0.9999	0.5740	0.4803	0.2091	0.1556	0.0034	0.0049
23	0.9999	0.9996	0.5672	0.4450	0.1846	0.1333	0.0036	0.0059
24	0.9999	0.9994	0.4894	0.3945	0.1443	0.1054	0.0056	0.0089
25	0.9997	0.9991	0.4625	0.3605	0.1194	0.0897	0.0087	0.0109
26	0.9993	0.9986	0.3721	0.3164	0.0825	0.0716	0.0161	0.0183
27	0.9995	0.9976	0.3233	0.2589	0.0710	0.0536	0.0227	0.0277



Figure 6: Power comparison: non-monotone hazard alternative at 10% when the sample size is 30.

5 Conclusion

As it is shown by Figures 4-6 and Tables 3-5 none of the available test statistics could beat others against all alternatives, but it is notable that the proposed test statistic shows better powers than the available test statistics against the alternatives with monotone increasing hazard functions, which applies to many real-life applications. For our test statistic against the alternatives monotone increasing hazard functions, Figures 4-6 show that with less remaining data, the proposed test statistic has very better performance than the other test statistics.

This work has the potential to be applied in the context of censored data and goodness of fit tests. This paper can elaborate further researches by extending such modifications for other censoring schemes such as progressive censoring scheme. Finally, this area of research can be expanded by considering other distributions besides the exponential distribution such as Pareto, Log normal and Weibull distributions.

References

[1] I. Arizono.I., Ohta.H. (1989) A test for normality based on Kullback-Leibler information, *American Statistician.*, 43, 20-23.

[2] S. Ascher.S. (1990) A survey of tests for exponentiality, *Comm. Statist. -Theor. Meth.*, 19, 1811-1825.

[3] Balakrishnan.N., Habibi Rad.A. and Arghami.N.R. (2007) Testing exponentiality based on Kullback-Leibler information with progressively type-II censored data, *IEEE Trans. Rel.*, 56, 349-356.

[4] Bowman.A.W. (1992) Density based tests for goodness-of-fit, *J. Statist. Comput. Simul.*, 40, 1-13.

[5] Brain.C.W., Shapiro.S.S. (1983) A regression test for exponentiality: censored, complete samples, *Technometrics.*, 25, 69-76.

[6] Dudewicz.E.J., Van Der Meulen.E.C. (1987) Emiric entropy, a new approach to non-parametric entropy estimation. In: Puir.M.L., Vilaplana.J.P. and Wrtz.W. (Eds.), New Perspectives in Theoretical and Applied Statistics, *Wiley, New York*, 202-207.

[7] Dudewicz.E.J., van der Meulen.E.C. (1981) Entropy-based tests of uniformity, *J. Amer. Statist. Assoc.*, 76, 967-974.

[8] Ebrahimi.N., Pflughoeft.K. and Soofi.E.S. (1994) Two measures of sample entropy, *Statist. Prob. Letters.*, 20, 225-234.

[9] Ebrahimi.N., Habibullah.M. and Soofi.E.S. (1992) Testing exponentiality based on Kullback-Leibler information, *J. Royal Statist. Soc.*, 54, 739-748.

[10] Ellinger.A.G. (1971) The Art of Investment, *Bowers and Bowers*, London.

[11] Gan.F.F., Koehler.K.J. (1990) Goodness-of-fit tests based on P-P probability plots, *Tecnometrics.*, 32, 289-303.

[12] Gokhale.D.V. (1983) On entropy-based goodness-of-fit tests, *Comput. Statist. Data Anal.*, 1, 157-165.

[13] Habibi Rad.A., Yousefzadeh.F. and Balakrishnan.N. (2011) Goodness of fit test based on Kullback-Leibler information for progressively type-II censored data, *IEEE Trans. Rel.*, 60, 570-579.

[14] Henze.N. (1993) A new flexible class of omnibus tests for exponentiality, *Comm. Statist. -Theor. Meth.*, 22, 115-133.

[15] Kallenberg.W.C.M., Ledwina.T. (1997) Data driven smooth tests for composite hypothesis: comparisons of powers, *J. Statist. Comput. Simul.*, 59, 101-121.

[16] Labie.E, private communication.

[17] LaRiccia.V. (1991) Smooth goodness of fit tests: a quantile function approach, *J. Amer. Statist. Assoc.*, 86, 427-431.

[18] Park.S. (2005) Testing exponentiality based on the KullbackLeibler information with the type II censored data, *IEEE Trans. Rel.*, 54, 22-26.

[19] Park.S. (1995) The entropy of consecutive order statistics, *IEEE Trans. Inform. Theory.*, 41, 2003-2007.

[20] Saaredra.A., Cao.R. (2000) On the estimation of the marginal density of a moving average process, *Canad. J. Statist.*, 28, 799-815.

[21] Samanta.M., Schwarz.C.J. (1988) The Shapiro-Wilk test for exponentiality based on censored data, *J. Amer. Statist. Assoc.*, 83, 528-531.

[22] Shannon.C.E. (1948) A mathematical theory of communications, *Bell System Tech. J.*, 27, 379-423.

[23] Silverman.B.W. (1986) Density Estimation for Statistics and Data Analysis, *Chapman and Hall*, London.

[24] Theil.H. (1980) The entropy of the maximum entropy distribution, *Economics Letters.*, 5, 145-148.

[25] Vasicek.O. (1976) A test for normality based on sample entropy, *J. Royal Statist. Soc.*, 38, 54-59.

[26] Wong.K.M., Chan.S. (1990) The entropy of ordered sequences and order statistics, *IEEE Trans. Inform. Theory.*, 36, 276-284.

[27] Yousefzadeh.F., Arghami.N.R. (2008) Testing exponentiality based on Type-II censored data and a new cdf estimator, *Comm. Statist. -Simul. Comput.*, 37, 1479-1499.


B1
B2
B3
B4
B5
B6
B7
B8
power
r
0
0.2
0.4
0.6
0.8
1
15
20
25
30
A
K